

# Red Cell Volume Expansion at Altitude: A Meta-analysis and Monte Carlo Simulation

PETER RASMUSSEN<sup>1,2,3</sup>, CHRISTOPH SIEBENMANN<sup>1,2</sup>, VÍCTOR DÍAZ<sup>1,4,5</sup>, and CARSTEN LUNDBY<sup>1,2</sup>

<sup>1</sup>Zurich Center of Integrative Human Physiology, Institute of Physiology, Zurich, SWITZERLAND; <sup>2</sup>Institute of Physiology, University of Zurich, Zurich, SWITZERLAND; <sup>3</sup>Department of Neuroscience and Pharmacology, University of Copenhagen, Copenhagen, DENMARK; <sup>4</sup>Institute of Veterinary Physiology, University of Zurich, Zurich, SWITZERLAND; <sup>5</sup>Department of Health and Human Performance, Faculty of Sport Science-INEF, Technical University of Madrid, Madrid, SPAIN

## ABSTRACT

RASMUSSEN, P., C. SIEBENMANN, V. DÍAZ, and C. LUNDBY. Red Cell Volume Expansion at Altitude: A Meta-analysis and Monte Carlo Simulation. *Med. Sci. Sports Exerc.*, Vol. 45, No. 9, pp. 1767–1772, 2013. **Introduction:** Altitude acclimatization is associated with a rapid increase in hematocrit. The time course and the contribution of the red cell volume expansion are not clear. The purpose of the present meta-analysis was to explore how much altitude exposure is required to induce polycythemia in healthy lowlanders. **Methods:** A systematic review was performed of 66 published articles (including 447 volunteers) identified through literature search. We performed a mixed-model random-effects meta-analysis and a Monte Carlo simulation on the extracted data. **Results:** The following results were obtained in this study: 1) the red cell volume expansion for a given duration of exposure is dependent on altitude ( $P < 0.0001$ ), that is, that the increase in red cell volume was accelerated at higher altitudes; and 2) the extent of the erythropoietic response depends on the initial red cell volume ( $P < 0.0001$ ). It seems that exposure time must exceed 2 wk at an altitude of more than 4000 m to exert a statistically significant effect. At lower altitudes, longer exposure times are needed with altitudes lower than 3000 m not yielding an increase within 4 wk. **Conclusions:** Red cell volume response to hypoxia is generally slow, although it accelerates with increasing altitude. This, in combination with a dependency on initial red cell volume, suggests that, for example, athletes may need to spend more time at altitude to see an effect on red cell volume than commonly recommended. **Key Words:** ALTITUDE TRAINING, HEMOGLOBIN MASS, HYPOXIA, SYSTEMATIC REVIEW

Hypoxia leads to the stabilization of the hypoxia inducible factor 2 system in the renal peritubular cells, which subsequently binds to the hypoxic response element on the erythropoietin (Epo) gene and induces synthesis and release of Epo (13). As a result, plasma Epo concentrations start to increase after approximately 2 h of hypoxic exposure (6). As exposure continues, a zenith is reached after 3–4 d, where after-plasma Epo concentration gradually decreases to stabilize values slightly above sea level (3,23,29).

The primary function of Epo is to regulate red blood cell volume by promoting the proliferation and differentiation of erythrocytic progenitors, and because permanent residence at altitude is associated with an augmented total red cell volume (RCV) (29), the paradigm has evolved that extended

hypoxic exposure induces polycythemia in lowlanders. Nevertheless, the Epo response to hypoxia does not necessarily translate into an elevated RCV (1,27). In fact, the initial increase in hematocrit at altitude is entirely the result of a plasma volume contraction that occurs within the first hours or days of exposure (26). In contrast, the time course of RCV expansion and the dependency on the degree of hypoxia remain largely unknown. This is related to the inconvenience of exposing a sufficient subject number to different altitudes for a prolonged time. To overcome this problem, we conducted a meta-analysis of published experiments and explored the altitude exposure and duration thereof required to induce polycythemia in healthy lowlanders.

## METHODS

**Search Strategy.** This meta-analysis is based on articles retrieved from the Web-based databases PubMed and Web of Science (see Supplementary Digital Content, <http://links.lww.com/MSS/A266>). Up to January 2012, we introduced the following key words and Boolean connectors: (altitude OR hypoxia) AND acclimatization AND (hemoglobin mass OR hemoglobin mass OR hematocrit OR hematocrit OR red blood cell mass OR red blood cell volume OR blood volume OR plasma volume).

The search was refined by applying the limits “Humans” and “All adult: +19 years” in PubMed and searching within the subject areas of physiology, respiratory system, cardiovascular system cardiology, sport sciences, and hematology in the Web of Science. This initial search generated a group of 6775 references that were transferred to an EndNote (Thomson Reuters, New York, NY) database.

In EndNote, duplicate references were removed. From the remaining references *in vitro*, animal, or high altitude natives studies, reviews and conference proceedings were eliminated. The abstracts of the 446 remaining articles were then independently evaluated by two researchers, and articles not fulfilling the eligibility criteria (see next section) were removed. Disagreements were resolved by involving the rest of the research team. By this process, 53 articles were selected.

Subsequently, we built a reference map for each article and obtained the references cited in or citing the initial 53 articles. This step retrieved a new group of 3185 articles (1503 backward and 1682 forward references), and the same process detailed earlier and in Figure 1 was repeated. Finally, 66 articles were selected to be included in the meta-analysis (see Supplementary Digital Content, <http://links.lww.com/MSS/A266>).

**Eligibility criteria.** Any study on healthy humans reporting data before and after a period of exposure to hypoxia (normobaric or hypobaric, continuous or intermittent) for blood compartments (plasma volume, RCV, or blood volume) or total hemoglobin mass was eligible. Only investigations using direct methods of measurement (Evans blue, carbon monoxide rebreathing, radioiodinated albumin, or brilliant red dilution) were accepted. Studies in which the experimental procedures included any intervention susceptible of affecting any blood compartment, other than physical activity, were eliminated as well.

**Data extraction.** The data were transferred into an Excel file by two researchers working independently and subsequently compared with avoid errors. When data were

extracted from the figures, the mean between the values obtained by the two researchers was taken.

When altitude was not constant, a time-weighted average was calculated.

$$\text{Average altitude} = \frac{\sum_{i=1}^n \text{Dur}_i \times \text{Alt}_i}{\text{Total duration}} = \frac{\text{Dur}_1 \times \text{Alt}_1 + \text{Dur}_2 \times \text{Alt}_2 + L + \text{Dur}_n \times \text{Alt}_n}{\text{Total duration}}$$


where  $\text{Dur}_i$  and  $\text{Alt}_i$  are the duration and altitude at step  $i$ . So for, for example, a stepwise ascent to Mount Everest (24), the average altitude is derived as  $[1 \text{ d} \times 3800 \text{ m} + 5 \text{ d} \times 4500 \text{ m} + 5 \text{ d} \times 5000 \text{ m} + 2 \text{ d} \times 5500 + \dots] / 36 \text{ d} = 5720 \text{ m}$ . If only SEM values were reported, variance was calculated from sample size and SD; otherwise, variance was calculated from SD alone.

**Statistical analysis.** A mixed-model random-effects meta-regression was performed (SAS 9.2; SAS Institute Inc., Cary, NC) on log-response ratios (12). The log-response ratio was calculated as the logarithm to the change in RCV, that is, the postvalue divided by the prevalue/control value. We chose RCV as our effect parameter because of the linear relationship with hemoglobin mass, whereas hematocrit and hemoglobin concentrations were excluded for their dependency on plasma volume. If RCV was not reported, it was derived from hemoglobin mass according to a linear relationship derived from the 13 studies ( $n = 28$ ) that reported both hemoglobin mass and RCV ( $\text{RCV} = \text{hemoglobin mass} \times 2.899$ ,  $P < 0.0001$ ). In general, none of the included studies reported confidence intervals or SD on the change in RCV; rather, SD (or SEM) was reported on the pre- and postvalues separately. Variance was therefore corrected as proposed for crossover trials (5).  $P$  values  $< 0.05$  were considered statistically significant.

## ANALYSIS AND RESULTS

Overall subject characteristics are reported in Table 1. In total, this analysis reports data from 447 volunteers of which 376 were men and 71 women. RCV increases versus exposure time and altitude are shown in Figure 1. Of the collected data, 75% documented an increase in RCV, that is, an effect size  $> 0$  regardless of statistical significance. Across the data set, average RCV increase per week was  $49 \pm 240 \text{ mL} \cdot \text{wk}^{-1}$ . Stratifying by exposure time, average RCV increases per week were  $37 \pm 81$ ,  $56 \pm 21$ , and  $100 \pm 467 \text{ mL} \cdot \text{wk}^{-1}$  with 10, 20, and 28 d of exposure, respectively, across all altitudes. For altitudes higher than 3000 m, the corresponding values were  $46 \pm 53$ ,  $46 \pm 258$ , and  $188 \pm 466 \text{ mL} \cdot \text{wk}^{-1}$ .

**Heterogeneity.** Because of the inclusion of articles spanning different methods (plasma dye dilution, carbon monoxide rebreathing, or radioactive albumin labeling), altitudes (from 1300 to 6000 m above sea level), exposure times (from 1 d to 1 yr), intermittent hypoxia (range  $1\text{--}21 \text{ h} \cdot \text{d}^{-1}$ , mean  $12 \pm 5 \text{ h} \cdot \text{d}^{-1}$ ), and normobaric hypoxia, data set heterogeneity is a potential confounder that might affect outcome of the analysis (Table 2). We tested for inhomogeneity



**FIGURE 1—Red blood cell response to exposure time and altitude.** Bubble sizes are scaled to log-response ratio and weighted according to study sample size to preserve presentation fidelity transformed with square root function. Filled bubbles indicate increased red cell increase, and open bubbles indicate decreases.

TABLE 1. Volunteer characteristics.


	5%	25%	Median	75%	95%
Hematocrit (%)	39.7	42.4	44.3	46.0	46.8
Body weight (kg)	56.5	58.9	66.1	72.6	74.1
[Hb] (mmol·L <sup>-1</sup> )	9.3	13.9	14.7	15.0	15.7
Hemoglobin mass (g)	654	733	868	988	1023
Per kg body weight	10.0	11.5	12.7	14.4	17.5
Plasma volume (mL)	2459	3019	3115	3536	4075
Per kg body weight	39.0	44.4	49.5	55.1	60.6
Red blood cell volume (mL)	1896	2125	2518	2864	2966
Per kg body weight	28.9	33.4	36.9	41.7	50.7
Total blood volume (mL)	4377	5410	5798	5979	7220
Per kg body weight	68.2	77.3	84.0	103.5	105.8

using a mixed-model regression analysis and found no significant effect of the measurement method ( $P = 0.94$ ), normobaric versus hypobaric hypoxia ( $P = 0.68$ ), continuous versus intermittent hypoxia ( $P = 0.14$ ), or exercise versus no exercise ( $P = 0.23$ ). Accordingly, all data were included in the main analysis.

**Effect of altitude and duration of exposure.** For clarity of presentation, data were divided into both altitude and duration quartiles (see Fig. 2). The random effects model yielded significant main effects for both altitude and duration (both  $P < 0.0001$ , Fig. 1); however, there was also a significant interaction effect ( $P < 0.0001$ ), and therefore the main effects ( $P$  values indicated) should be interpreted with caution. It seems that exposure time must exceed at least 2 wk at an altitude of more than 4000 m to exert a significant effect. At lower altitudes, even longer exposure times are needed with altitudes lower than 3000 m not yielding a statistically significant result within 4 wk.

We subsequently performed Monte Carlo simulation to estimate the required exposure time for an increase in RCV at different altitudes. We entered mean pre- and postexposure RCV values with SD along with exposure time into a custom-written Matlab (MathWorks, Natick, MA) procedure. From these distributions, we randomly drew paired numbers and calculated changes in RCV as a function of time. The results of the simulation are presented in Table 3. There is an accelerating effect of increasing altitude so that, for example, reaching a 10% increase at 2000 m will take 42 to 145 d, whereas at 3500 m this can be achieved between 23 and 51 d.

**Effect of initial RCV.** Finally, we examined the effect of the initial RCV on the RCV response to hypoxic exposure. To obtain a homologous data set, that is, data obtained with comparable exposure and methodology, we restricted the analysis to the special case of exposure between 3000 and



**FIGURE 2—Red blood cell response.** Data are presented in altitude and exposure quartiles. Note that there was a significant interaction between the effect of altitude and the exposure duration ( $P < 0.0001$ ); therefore, the main effects ( $P$  values indicated) should be interpreted with caution. It is, however, clear that exposure time must exceed 17 d at an altitude of more than 4000 m to exert a significant effect on the oxygen carrying capacity of the blood. Data are weighted log-response ratios with error bars indicating 95% confidence intervals.

3500 m and more than 7 d of stay. The random-effects model showed a highly significant effect of initial RCV with a high initial RCV showing less increase compared with low initial RCV ( $P < 0.0001$ , Fig. 3).

## DISCUSSION

The main findings of the meta-analysis and Monte Carlo simulations are that altitude exposure must exceed 2 wk at an altitude of  $>4000$  m to exert a significant effect on RCV. At lower altitudes, even longer exposure is required with altitudes lower than 3000 m not yielding a statistically significant result within 4 wk. Thus, we established an altitude dose-response curve, indicating that the hypoxia-induced response in RCV may be slower than widely expected (30).

**Variation in RCV response.** We found that the average RCV increase per week for our data was  $49 \pm 240$  mL·wk<sup>-1</sup>, which, on average, is in accordance with previous findings with EPO administration (2). Although we find the large variation surprising, similar results may be found in the plasma Epo response to altitude. Ge et al. (10) reported that the increase in plasma Epo after 24 h at 2800 m varied markedly among individuals ( $n = 48$ ), ranging from

TABLE 2. No. data points entering the analysis depending on methodology.

	<i>N</i>	Pct.	CO Rebreathing	Plasma Dye Dilution	Radiolabeled Albumin	Hypobaric Hypoxia	Normobaric Hypoxia	Continuous Hypoxia	Intermittent Hypoxia
CO rebreathing	67	44.1	—	—	—	—	—	—	—
Plasma dye dilution	56	36.8	—	—	—	—	—	—	—
Radiolabeled albumin	29	19.1	—	—	—	—	—	—	—
Hypobaric hypoxia	120	79.0	35	56	29	—	—	—	—
Normobaric hypoxia	32	21.1	32	0	0	—	—	—	—
Continuous hypoxia	111	73.0	31	52	28	111	0	—	—
Intermittent hypoxia	41	27.0	36	4	1	9	32	—	—
No exercise	92	60.5	22	52	18	92	0	91	1
Exercise	60	39.5	45	4	11	28	32	20	40



TABLE 3. Exposure time in days to obtain 95% probability for an increase in red blood cell volume.

Both Continuous and Intermittent Hypoxia	Altitude			
	2000	2500	3000	3500
1.00%	27 (21–39)	22 (18–32)	24 (18–31)	20 (12–38)
2.50%	35 (27–46)	25 (20–36)	26 (21–32)	20 (13–40)
5.00%	45 (33–60)	32 (28–44)	30 (26–38)	22 (17–41)
7.50%	59 (37–94)	42 (35–56)	33 (28–42)	23 (21–47)
10.00%	85 (42–145)	49 (39–116)	37 (31–46)	26 (23–51)
Only continuous hypoxia				
1.00%	30 (24–44)	18 (14–23)	21 (13–28)	20 (12–40)
2.50%	38 (31–48)	20 (16–26)	21 (15–28)	21 (13–38)
5.00%	52 (37–72)	29 (22–34)	23 (19–30)	22 (17–41)
7.50%	69 (40–99)	34 (25–47)	24 (21–33)	23 (21–45)
10.00%	95 (47–141)	37 (28–49)	26 (23–37)	26 (23–52)

Monte Carlo simulation, 50,000 simulations; each altitude spans 375 m on each side of the midpoint. Numbers in parentheses are the 5th–95th percentile range.

–41% to 400%, which seemed mainly governed by upstream factors related to renal parenchymal PO<sub>2</sub>, although genetic factors cannot be ruled out (19). Although variations in both RCV and the Epo response to altitude are observed, these do not seem to be interlinked. In a retrospectively analysis (16), no correlation between EPO and erythropoietic response was observed, which points to interindividual variation in responsiveness (4). The Epo-induced expansion of RCV can be used as a measuring stick for the speed of which RCV expands at altitude. Previous findings with Epo suggest that RCV expansion would not be expected to exceed 50 mL·wk<sup>–1</sup> (2); however, we found that >35% of the reported data correspond to RCV increases per week in excess of 50 mL·wk<sup>–1</sup>, and this finding is in accordance with Sawka et al. (26). It is intriguing that, in some studies, an increase in RCV is observed within days of exposure to an altitude that did not stimulate polycythemia even after weeks in other studies (Fig. 1). This suggests a substantial interindividual variation (27). Other candidates that could lead to variation include the use of different measurement techniques or of the application of continuous versus intermittent or normobaric versus hypobaric hypoxia (Table 2). Although there was no systematic effect observed when these parameters were entered as factors, the inclusion of results obtained with different methods may have increased the variance and reduced the power of the analysis. Some evidence suggests slightly different physiological responses to hypobaric versus normobaric hypoxia about ventilation and acute mountain sickness (18,22). However, this does not seem to be the case for the RCV response, which is in line with the observation that the Epo response to hypobaric and normobaric hypoxia is also similar (7).

**Effect of initial RCV.** One factor that affected the response to altitude exposure was the initial RCV. In this context, we recently failed to observe an increase in RCV after 16 h·d<sup>–1</sup> at 3000 m for 4 wk (27). From the analysis, we would expect a 5%–10% increase in RCV, and we speculated (25) that the absence of an increase in RCV could be related to the high initial hemoglobin mass of the included elite athletes that were at the upper end of physiological range (Table 1). The present meta-analysis supports this approach (Fig. 3). Although it is intuitively straightforward that an upper limit to RCV expansion must exist, the physiological

mechanism governing such a limit is not as easily explained. Although the current meta-analysis indicates that a high initial RCV limits a further expansion in hypoxia, the study by Kapoor and Chatterjee (15) pointed out that despite a high baseline level, an increase in RCV can still occur if duration and altitude are sufficient (Fig. 1). Thus, to what extent a physiological upper limit to RCV exists is not clear from this meta-analysis. However, 17.5 g of hemoglobin per kilogram bodyweight has been reported in World and Olympic Champion cross-country skiers (14), which is markedly higher than the values reported in healthy but untrained high altitude Peruvians (8). Thus, the upper limit is likely not reached by altitude exposure alone. It should thus be considered that an RCV ceiling may not relate to increased oxygen carrying capacity of blood alone. Any such ceiling could also be related to an upper limit in blood volume and thereby to blood pressure regulation.

Another explanation for the finding that a high initial RCV attenuates the further increase with hypoxic exposure could be the statistical phenomenon of regression toward the mean. Regression toward the mean may occur when RCV is




FIGURE 3—The effect of initial hemoglobin mass on response to hypoxic exposure. To ensure clarity of presentation and to avoid confounding bias, data are limited to exposure times between 7 and 40 d at altitudes between 3000 and 3500 m. On the abscissae, log-response ratios with 95% confidence intervals are presented and on the ordinate hemoglobin mass with SD. Mean values for altitude (3327, 3097, and 3070 m) and exposure (21, 19, and 23 d) for the three hemoglobin mass groups (*N* = 13, 14, and 15) are not different.

high on its first measurement because it will by chance tend to be closer to the average on a second measurement. To avoid drawing wrong conclusions from this meta-analysis (particularly Fig. 3), the possibility of regression toward the mean should be considered. However, because regression toward the mean is not based on cause and effect, but rather on random error in a natural distribution around a mean, we cannot exclude any underlying physiological mechanisms influencing the effect of initial RCV on erythropoietic response to altitude exposure.

**Is altitude exposure relevant for athletes?** To increase RCV and thereby also sea-level exercise performance, it is recommended that athletes reside in normobaric or hypobaric hypoxia corresponding to an altitude of 2500–3000 m for a minimum of 14 h·d<sup>-1</sup> for 3 wk (30). The effect of the initial RCV (Fig. 3) and the considerable time required to establish an RCV response to medium altitude exposure (Table 3) are of relevance for athletes considering engaging into altitude training. Assuming that anecdotal reports of athletes coping poorly with altitudes higher than 3000 m (increased recovery period after exercise, poor quality of sleep, etc.) are correct, then altitudes above this threshold should be avoided. Because only studies in athletes who have demonstrated >5% increase in hemoglobin mass after altitude training generally report an increase in exercise performance (25), it is recommended that athletes should spend sufficient time at altitude to achieve a corresponding increase in hemoglobin mass. A gain in hemoglobin mass of 7.5% requires between 35 and 56 d of live high–train low or 25 to 47 d of continuous exposure at 2500 m to achieve a >95% probability for an increase (Table 3). This may be further extended by a high initial RCV, which results in lower probability for a further increase to occur. This questions the feasibility of altitude training for elite athletes, and the potential gains by altitude training, at least in athletes with a high RCV starting point, should therefore be carefully reconsidered.

**Publication bias.** We found some indications of publication bias in the tendency for studies with small changes and large variations to be missing from the analysis. Thus, we cannot exclude that the analysis overestimates the positive outcome. On the other hand, almost 40% of the included data points reported negative findings and 50% of the data points reported less than 2.5% increase.

**Data set heterogeneity.** Roughly a fourth of the data originates from studies involving intermittent hypoxia (Table 2). We were not able to detect any significant differences when the two hypoxic exposures were compared in the random-effects model. However, Monte Carlo analysis revealed that when the data obtained from intermittent hypoxic exposure were removed from the analysis, time to a 95% chance for an increase in RCV generally decreased with 4–7 d, particularly around 3000 m above sea level where most of the intermittent hypoxia studies were performed. We also recorded iron supplementation, and in 72% of the studies, no iron supplementation was performed. Levine and Stray-Gundersen (17) suggested that iron

supplementation is needed only if iron stores are low. Parisotto et al. (20) could not find an effect of iron supplementation when Epo was administered. Thus, iron availability may not be an issue for healthy individuals. However, it is still not straightforward to assess the impact of iron supplementation as, for example, most studies with iron supplementation use intermittent hypoxia. Thus, there is a risk of covariance between the parameters. This extends to possible differences between normobaric and hypobaric hypoxia, sex, and methods used. In general, we saw no indications that normobaric hypoxia was producing a different response than hypobaric hypoxia, which is in accordance with our recent findings (21). Likewise, we were not able to detect any differences between the few studies reporting only women and those with only men or a mixed population. What we note, however, is that to our knowledge, no studies have looked specifically at sex differences in RCV response to altitude dwelling, and very few studies report males and females separately. Finally, the method to estimate RCV may also introduce bias. According to Sawka et al. (26), CO rebreathing may overestimate not only the RCV but also the changes in RCV after altitude exposure because of CO loss to myoglobin or other iron porphyrin molecules (i.e., in the liver) and according changes in the quantity of these molecules after altitude exposure. However, Gore et al. (11) and Thomsen et al. (28) concluded that the two measurements were equivalent. Furthermore, CO loss to myoglobin is assumed minimal (1.6 mL for 10 min [9]), and even if myoglobin should change because of altitude exposure, the effect hereof on RCV must be assumed neglectable. Finally, it is unknown if these extravascular iron molecules actually increase with hypoxic exposure.

## CONCLUSIONS

The physiological process of red cell expansion occurs relatively slowly with only minor chances for an increase to occur within 2 wk and exposure times longer than 4 wk generally required. Also, the accepted paradigm that altitudes <3000 m are sufficient to trigger a robust RCV should be reconsidered, particularly if the initial RCV is high. This, in combination with a dependence on initial RCV, suggests that, for example, athletes may need to spend more time at altitude to see an effect on RCV than commonly recommended.

This study was supported by the Danish National Research Council (grant no. 10-094110) and the Zurich Center of Integrative Human Physiology.

Peter Rasmussen, Christoph Siebenmann, and Víctor Díaz contributed equally to the manuscript.

C.L. initiated the investigation; P.R. and V.D. designed the literature search; C.S. and V.D. extracted the data; and P.R. performed the analysis. All authors wrote the article and approved the final version.


The authors declare no conflict of interest.

The results of the present study do not constitute endorsement by the American College of Sports Medicine.

## REFERENCES

- Ashenden MJ, Gore CJ, Martin DT, Dobson GP, Hahn AG. Effects of a 12-day “live high, train low” camp on reticulocyte production and haemoglobin mass in elite female road cyclists. *Eur J Appl Physiol.* 1999;80:472–8.
- Berglund B, Ekblom B. Effect of recombinant human erythropoietin treatment on blood pressure and some haematological parameters in healthy men. *J Intern Med.* 1991;229(2):125–30.
- Berglund B, Gennser M, Örnhaugen H, Östberg C, Wide L. Erythropoietin concentrations during 10 days of normobaric hypoxia under controlled environmental circumstances. *Acta Physiol Scand.* 2002;174(3):225–9.
- Chapman RF, Stray-Gundersen J, Levine BD. Individual variation in response to altitude training. *J Appl Physiol.* 1998;85(4):1448–56.
- Curtin F, Altman DG, Elbourne D. Meta-analysis combining parallel and cross-over clinical trials. I: continuous outcomes. *Stat Med.* 2002;21(15):2131–44.
- Eckardt KU, Boutellier U, Kurtz A, Schopen M, Koller EA, Bauer C. Rate of erythropoietin formation in humans in response to acute hypobaric hypoxia. *J Appl Physiol.* 1989;66(4):1785–8.
- Fandrey J. Oxygen-dependent and tissue-specific regulation of erythropoietin gene expression. *Am J Physiol Regul Integr Comp Physiol.* 2004;286(6):R977–88.
- Gamboa A, Gamboa JL, Holmes C, et al. Plasma catecholamines and blood volume in native Andeans during hypoxia and normoxia. *Clin Auton Res.* 2006;16(1):40–5.
- Garvican LA, Burge CM, Cox AJ, Clark SA, Martin DT, Gore CJ. Carbon monoxide uptake kinetics of arterial, venous and capillary blood during CO rebreathing. *Exp Physiol.* 2010;95(12):1156–66.
- Ge RL, Witkowski S, Zhang Y, et al. Determinants of erythropoietin release in response to short-term hypobaric hypoxia. *J Appl Physiol.* 2002;92(6):2361–7.
- Gore CJ, Hopkins WG, Burge CM. Errors of measurement for blood volume parameters: a meta-analysis. *J Appl Physiol.* 2005;99(5):1745–58.
- Hedges LV, Gurevitch J, Curtis PS. The meta-analysis of response ratios in experimental ecology. *Ecology.* 1999;80(4):1150–6.
- Jelkmann W. Erythropoietin after a century of research: younger than ever. *Eur J Haematol.* 2007;78(3):183–205.
- Jelkmann W. Regulation of erythropoietin production. *J Physiol.* 2011;589(Pt 6):1251–8.
- Kapoor SC, Chatterjee AK. Hematological response among new arrival at high-altitude. *Indian J Med Res.* 1978;67(Mar):428–34.
- Levine BD, Stray-Gundersen J. “Living high-training low”: effect of moderate-altitude acclimatization with low-altitude training on performance. *J Appl Physiol.* 1997;83(1):102–12.
- Levine BD, Stray-Gundersen J. A practical approach to altitude training: where to live and train for optimal performance enhancement. *Int J Sports Med.* 1992;13(1 Suppl):S209–12.
- Loeppky JA, Icenogle M, Scotto P, Robergs R, Hinghofer-Szalkay H, Roach RC. Ventilation during simulated altitude, normobaric hypoxia and normoxic hypobaria. *Respir Physiol.* 1997;107(3):231–9.
- Ou LC, Salceda S, Schuster SJ, et al. Polycythemic responses to hypoxia: molecular and genetic mechanisms of chronic mountain sickness. *J Appl Physiol.* 1998;84(4):1242–51.
- Parisotto R, Wu M, Ashenden MJ, et al. Detection of recombinant human erythropoietin abuse in athletes utilizing markers of altered erythropoiesis. *Haematologica.* 2001;86(2):128–37.
- Rasmussen P, Nordsborg N, Taudorf S, et al. Brain and skin do not contribute to the systemic rise in erythropoietin during acute hypoxia in humans. *FASEB J.* 2012;26(5):1831–4.
- Roach RC, Loeppky JA, Icenogle MV. Acute mountain sickness: increased severity during simulated altitude compared with normobaric hypoxia. *J Appl Physiol.* 1996;81(5):1908–10.
- Robach P, Cairo G, Gelfi C, et al. Strong iron demand during hypoxia-induced erythropoiesis is associated with down-regulation of iron-related proteins and myoglobin in human skeletal muscle. *Blood.* 2007;109(11):4724–31.
- Robach P, Dechaux M, Jarrot S, et al. Operation Everest III: role of plasma volume expansion on VO<sub>2</sub>(max) during prolonged high-altitude exposure. *J Appl Physiol.* 2000;89(1):29–37.
- Robach P, Lundby C. Is live high—train low altitude training relevant for elite athletes with already high total hemoglobin mass? *Scand J Med Sci Sports.* 2012;22(3):303–305.
- Sawka MN, Convertino VA, Eichner ER, Schnieder SM, Young AJ. Blood volume: importance and adaptations to exercise training, environmental stresses, and trauma/sickness. *Med Sci Sports Exerc.* 2000;32(2):332–48.
- Siebenmann C, Robach P, Jacobs RA, et al. “Live high—train low” using normobaric hypoxia: a double-blinded, placebo-controlled study. *J Appl Physiol.* 2012;112(1):106–17.
- Thomsen JK, Fogh-Andersen N, Bulow K, Devantier A. Blood and plasma volumes determined by carbon monoxide gas, 99mTc-labelled erythrocytes, 125I-albumin and the T 1824 technique. *Scand J Clin Lab Invest.* 1991;51(2):185–90.
- Weil JV, Jamieson G, Brown DW, Grover RF. The red cell mass-arterial oxygen relationship in normal man: application to patients with chronic obstructive airway disease. *J Clin Invest.* 1968;47(7):1627–39.
- Wilber RL, Stray-Gundersen J, Levine BD. Effect of hypoxic “dose” on physiological responses and sea-level performance. *Med Sci Sports Exerc.* 2007;39(9):1590–9.

## Supplemental Digital Content



**Figure.** Schematic view of the literature review process. Papers were initially retrieved from Web of Science and PubMed.

## Included references

- Alexander JK, Hartley LH, Modelski M, Grover RF. Reduction of stroke volume during exercise in man following ascent to 3,100 m altitude. *J Appl Physiol.* 1967;23(6):849-858. Prepublished on 1967/12/01 as DOI.
- Ashenden MJ, Gore CJ, Dobson GP, Hahn AG. "Live high, train low" does not change the total haemoglobin mass of male endurance athletes sleeping at a simulated altitude of 3000 m for 23 nights. *Eur J Appl Physiol Occup Physiol.* 1999;80(5):479-484. 10.1007/s004210050621.
- Ashenden MJ, Gore CJ, Martin DT, Dobson GP, Hahn AG. Effects of a 12-day "live high, train low" camp on reticulocyte production and haemoglobin mass in elite female road cyclists. *Eur J Appl Physiol.* 1999;80:472-478.
- Balke B, Nagle FJ, Daniels J. Altitude and Maximum Performance in Work and Sports Activity. *JAMA.* 1965;194(6):646-&.

5. Boning D, Maassen N, Jochum F, et al. After-effects of a high altitude expedition on blood. *Int J Sports Med.* 1997;18(3):179-185. Prepublished on 1997/04/01 as DOI 10.1055/s-2007-972616 [doi].
6. Brugniaux JV, Schmitt L, Robach P, et al. Eighteen days of "living high, training low" stimulate erythropoiesis and enhance aerobic performance in elite middle-distance runners. *J Appl Physiol.* 2006;100(1):203-211.
7. Clark SA, Quod MJ, Clark MA, Martin DT, Saunders PU, Gore CJ. Time course of haemoglobin mass during 21 days live high:train low simulated altitude. *Eur J Appl Physiol.* 2009;106(3):399-406. 10.1007/s00421-009-1027-4.
8. Clinton M, Thorn GW, Davenport VD. Studies on Altitude Tolerance .2. Studies on Normal Human Subjects - Effect of Repeated Short Exposures to Reduced Atmospheric Pressure. *Bulletin of the Johns Hopkins Hospital.* 1946;79(1):70-89.
9. Dehnert C, Hutler M, Liu Y, et al. Erythropoiesis and performance after two weeks of living high and training low in well trained triathletes. *Int J Sports Med.* 2002;23(8):561-566. Prepublished on 2002/11/20 as DOI 10.1055/s-2002-35533 [doi].
10. Dempsey JA, Reddan WG, Birnbaum ML, et al. Effects of acute through life-long hypoxic exposure on exercise pulmonary gas exchange. *Respir Physiol.* 1971;13(1):62-89. 10.1016/0034-5687(71)90065-x.
11. Dill DB, Braithwaite K, Adams WC, Bernauer EM. Blood-Volume of Middle-Distance Runners - Effect of 2,300-M Altitude and Comparison with Non-Athletes. *Med Sci Sports Exerc.* 1974;6(1):1-7.
12. Dill DB, Horvath SM, Dahms TE, Parker RE, Lynch JR. Hemoconcentration at altitude. *J Appl Physiol.* 1969;27(4):514-518.
13. Douglas CG, Halpane JS, Henderson Y, Schneider EC. Physiological observations made on Pike's Peak, Colorado, with special reference to adaptation to low barometric pressures. *Proc R Soc Lond B Biol Sci.* 1913;203:185-318. 10.1098/rstb.1913.0006.
14. Faura J, Reynafar C. Use metha-androstenolone for acceleration of erythropoiesis in exposure to height. *Arch Inst Biol Andina.* 1970;3(3-4):94-98.
15. Frayser R, Rennie ID, Gray GW, Houston CS. Hormonal and electrolyte response to exposure to 17,500 ft. *J Appl Physiol.* 1975;38(4):636-642.
16. Frese F, Friedmann-Bette B. Effects of Repetitive Training at Low Altitude on Erythropoiesis in 400 and 800 m Runners. *Int J Sports Med.* 2010;31(6):382-388.
17. Friedmann B, Frese F, Menold E, Bartsch P. Individual variation in the reduction of heart rate and performance at lactate thresholds in acute normobaric hypoxia. *Int J Sports Med.* 2005;26(7):531-536. 10.1055/s-2004-821326.



18. Friedmann B, Jost J, Rating T, et al. Effects of Iron Supplementation on Total Body Hemoglobin During Endurance Training at Moderate Altitude. *Int J Sports Med*. 1999;20(02):78,85. 10.1055/s-2007-971097.
19. Garvican LA, Pottgiesser T, Martin DT, Schumacher YO, Barras M, Gore CJ. The contribution of haemoglobin mass to increases in cycling performance induced by simulated LHTL. *Eur J Appl Physiol*. 2011;111(6):1089-1101. Prepublished on 2010/11/30 as DOI 10.1007/s00421-010-1732-z.
20. Gore CJ, Hahn A, Rice A, et al. Altitude training at 2690m does not increase total haemoglobin mass or sea level VO<sub>2</sub>max in world champion track cyclists. *J Sci Med Sport*. 1998;1(3):156-170. Prepublished on 1998/10/23 as DOI.
21. Gore CJ, Hahn AG, Burge CM, Telford RD. VO<sub>2</sub>max and haemoglobin mass of trained athletes during high intensity training. *Int J Sports Med*. 1997;18(6):477-482. Prepublished on 1997/08/01 as DOI 10.1055/s-2007-972667 [doi].
22. Gore CJ, Rodriguez FA, Truijens MJ, Townsend NE, Stray-Gundersen J, Levine BD. Increased serum erythropoietin but not red cell production after 4 wk of intermittent hypobaric hypoxia (4,000-5,500 m). *J Appl Physiol*. 2006;101(5):1386-1393. 10.1152/japplphysiol.00342.2006.
23. Greenleaf JE, Bernauer EM, Adams WC, Juhos L. Fluid-Electrolyte Shifts and Vo 2 Max in Man at Simulated Altitude (2,287 M). *J Appl Physiol*. 1978;44(5):652-658.
24. Grover RF, Selland MA, McCullough RG, et al. beta-adrenergic blockade does not prevent polycythemia or decrease in plasma volume in men at 4300 m altitude. *Eur J Appl Physiol*. 1998;77(3):264-270.
25. Hannon JP, Shields JL, Harris CW. Effects of Altitude Acclimatization on Blood Composition of Women. *J Appl Physiol*. 1969;26(5):540-547.
26. Hannon JP, Vogel JA. Oxygen transport during early altitude acclimatization: a perspective study. *Eur J Appl Physiol Occup Physiol*. 1977;36(4):285-297. Prepublished on 1977/05/10 as DOI.
27. Hansen JM, Olsen NV, Feldt-Rasmussen B, et al. Albuminuria and overall capillary permeability of albumin in acute altitude hypoxia. *J Appl Physiol*. 1994;76(5):1922-1927. Prepublished on 1994/05/01 as DOI.
28. Heinicke K, Heinicke I, Schmidt W, Wolfarth B. A three-week traditional altitude training increases hemoglobin mass and red cell volume in elite biathlon athletes. *Int J Sports Med*. 2005;26(5):350-355. 10.1055/s-2004-821052.
29. Imoberdorf R, Garlick PJ, McNurlan MA, et al. Enhanced synthesis of albumin and fibrinogen at high altitude. *J Appl Physiol*. 2001;90(2):528-537.

30. Jain SC, Bardhan J, Swamy YV, Krishna B, Nayar HS. Body-Fluid Compartments in Humans during Acute High-Altitude Exposure. *Aviat Space Environ Med.* 1980;51(3):234-236.
31. Jung RC, Dill DB, Horton R, Horvath SM. Effects of age on plasma aldosterone levels and hemoconcentration at altitude. *J Appl Physiol.* 1971;31(4):593-&.
32. Kapoor SC, Chatterjee AK. Hematological response among new arrival at high-altitude. *Indian J Med Res.* 1978;67(MAR):428-434.
33. Krzywicki HJ, Consolazio CF, Matoush LO, Johnson HL, Barnhart RA. Body composition changes during exposure to altitude. *Fed Proc.* 1969;28(3):1190-&.
34. Levine BD, Stray-Gundersen J. "Living high-training low": effect of moderate-altitude acclimatization with low-altitude training on performance. *J Appl Physiol.* 1997;83(1):102-112. Prepublished on 1997/07/01 as DOI.
35. Merino CF. Studies on Blood Formation and Destruction in the Polycythemia of High Altitude. *Blood.* 1950;5(1):1-31.
36. Neya M, Enoki T, Kumai Y, Sugoh T, Kawahara T. The effects of nightly normobaric hypoxia and high intensity training under intermittent normobaric hypoxia on running economy and hemoglobin mass. *J Appl Physiol.* 2007;103(3):828-834. 10.1152/japplphysiol.00265.2007.
37. Parving HH. The effect of hypoxia and carbon monoxide exposure on plasma volume and capillary permeability to albumin. *Scand J Clin Lab Invest.* 1972;30(1):49-56. 10.3109/00365517209081089.
38. Picon-Reategui E, Buskirk ER, Baker PT. Blood glucose in high-altitude natives and during acclimatization to altitude. *J Appl Physiol.* 1970;29(5):560-563. Prepublished on 1970/11/01 as DOI.
39. Pottgiesser T, Ahlgrim C, Ruthardt S, Dickhuth H-H, Schumacher YO. Hemoglobin mass after 21 days of conventional altitude training at 1816 m. *J Sci Med Sport.* 2009;12(6):673-675. 10.1016/j.jsams.2008.06.005.
40. Poulsen TD, Klausen T, Richalet JP, Kanstrup IL, Fogh-Andersen N, Olsen NV. Plasma volume in acute hypoxia: comparison of a carbon monoxide rebreathing method and dye dilution with Evans' blue. *Eur J Appl Physiol Occup Physiol.* 1998;77(5):457-461. Prepublished on 1998/04/30 as DOI.
41. Pugh LGC. Blood Volume and Haemoglobin Concentration at Altitudes above 18000 Ft (5500 M). *J Physiol.* 1964;170(2):344-354.
42. Reynafarje C, Lozano R, Valdivieso J. The polycythemia of high altitudes - iron metabolism and related aspects. *Blood.* 1959;14(4):433-455.

43. Robach P, Dechaux M, Jarrot S, et al. Operation Everest III: role of plasma volume expansion on VO<sub>2</sub>(max) during prolonged high-altitude exposure. *J Appl Physiol*. 2000;89(1):29-37. Prepublished on 2000/07/25 as DOI.
44. Robach P, Lafforgue E, Olsen NV, et al. Recovery of plasma volume after 1 week of exposure at 4,350 m. *Pflugers Arch*. 2002;444(6):821-828.
45. Robach P, Schmitt L, Brugniaux JV, et al. Living high-training low: effect on erythropoiesis and maximal aerobic performance in elite Nordic skiers. *Eur J Appl Physiol*. 2006;97(6):695-705.
46. Robach P, Schmitt L, Brugniaux JV, et al. Living high-training low: effect on erythropoiesis and aerobic performance in highly-trained swimmers. *Eur J Appl Physiol*. 2006;96(4):423-433.
47. Robertson EY, Aughey RJ, Anson JM, Hopkins WG, Pyne DB. Effects of Simulated and Real Altitude Exposure in Elite Swimmers. *J Strength Cond Res*. 2010;24(2):487-493.
48. Robertson EY, Saunders PU, Pyne DB, Aughey RJ, Anson JM, Gore CJ. Reproducibility of performance changes to simulated live high/train low altitude. *Med Sci Sports Exerc*. 2010;42(2):394-401. Prepublished on 2009/11/21 as DOI 10.1249/MSS.0b013e3181b34b57 [doi].
49. Robertson EY, Saunders PU, Pyne DB, Gore CJ, Anson JM. Effectiveness of intermittent training in hypoxia combined with live high/train low. *Eur J Appl Physiol*. 2010;110(2):379-387. 10.1007/s00421-010-1516-5.
50. Saunders PU, Ahlgrim C, Vallance B, et al. An Attempt to Quantify the Placebo Effect From a Three-Week Simulated Altitude Training Camp in Elite Race Walkers. *Int J Sports Physiol Perform*. 2010;5(4):521-534.
51. Saunders PU, Telford RD, Pyne DB, et al. Improved running economy in elite runners after 20 days of simulated moderate-altitude exposure. *J Appl Physiol*. 2004;96(3):931-937. 10.1152/jappphysiol.00725.2003.
52. Saunders PU, Telford RD, Pyne DB, Hahn AG, Gore CJ. Improved running economy and increased hemoglobin mass in elite runners after extended moderate altitude exposure. *J Sci Med Sport*. 2009;12(1):67-72. 10.1016/j.jsams.2007.08.014.
53. Siebenmann C, Robach P, Jacobs RA, et al. "Live high-train low" using normobaric hypoxia: a double-blinded, placebo-controlled study. *J Appl Physiol*. 2012;112(1):106-117. Prepublished on 2011/10/29 as DOI jappphysiol.00388.2011 [pii] 10.1152/jappphysiol.00388.2011.

54. Siggaard-Andersen J, Petersen FB, Hansen TI, Møllemaard K. Plasma volume and vascular permeability during hypoxia and carbon monoxide exposure. *Scand J Clin Lab Invest Suppl.* 1968;103:39-48. Prepublished on 1968/01/01 as DOI.
55. Singh MV, Jain SC, Rawal SB, et al. Comparative-study of acetazolamide and spironolactone on body-fluid compartments on induction to high-altitude. *Int J Biometeorol.* 1986;30(1):33-41. 10.1007/bf02192056.
56. Smith HP, Belt AE, Arnold HR, Carrier EB. Blood volume changes at high altitude. *Am J Physiol.* 1925;71(2):395-412.
57. Stokke KT, Rootwelt K, Wergeland R, Vale JR. Changes in plasma and red-cell volumes during exposure to high-altitude. *Scand J Clin Lab Invest.* 1986;46:113-117.
58. Surks MI, Chinn KS, Matoush LR. Alterations in body composition in man after acute exposure to high altitude. *J Appl Physiol.* 1966;21(6):1741-1746. Prepublished on 1966/11/01 as DOI.
59. Svedenhag J, Piehl-Aulin K, Skog C, Saltin B. Increased left ventricular muscle mass after long-term altitude training in athletes. *Acta Physiol Scand.* 1997;161(1):63-70. Prepublished on 1997/10/06 as DOI.
60. Takeno Y, Kamijo YI, Nose H. Thermoregulatory and aerobic changes after endurance training in a hypobaric hypoxic and warm environment. *J Appl Physiol.* 2001;91(4):1520-1528. Prepublished on 2001/09/25 as DOI.
61. Terzioglu M, Tuna N. Variations in Blood Volume at 1.85 Km Altitude. *J Appl Physiol.* 1954;6(7):417-422.
62. Turner HS, Hoffler GW, Billings CE, Bason R. An attempt to produce acclimatization to hypoxia by intermittent altitude exposure with vigorous exercise. *Aerosp Med.* 1969;40(9):971-976. Prepublished on 1969/09/01 as DOI.
63. Wehrli JP, Marti B. Live high-train low associated with increased haemoglobin mass as preparation for the 2003 World Championships in two native European world class runners. *Br J Sports Med.* 2006;40(2). e3  
10.1136/bjsm.2005.019729.
64. Wehrli JP, Zuest P, Hallen J, Marti B. Live high-train low for 24 days increases hemoglobin mass and red cell volume in elite endurance athletes. *J Appl Physiol.* 2006;100(6):1938-1945. 10.1152/jappphysiol.01284.2005.
65. Whitten BK, Burlingt.Rf, Posiviat.Ma, Sidel CM, Beecher GR. Amino Acid Catabolism in Environmental Extremes - Effect of High Altitude and Calories. *Am J Physiol.* 1970;218(5):1346-1350.



66. Wolfel EE, Groves BM, Brooks GA, et al. Oxygen transport during steady-state submaximal exercise in chronic hypoxia. *J Appl Physiol.* 1991;70(3):1129-1136. Prepublished on 1991/03/01 as DOI.